
L Number	Hits	Search Text	DB	Time stamp
1	0	mRNA near5 RT-PCR and circular adj1 cDNA	USPAT;	2003/02/11 12:38
		-	US-PGPUB;	
			DERWENT	
2	1697	mRNA near5 RT-PCR	USPAT;	2003/02/11 12:38
			US-PGPUB;	
			DERWENT	
3	0	(mRNA near5 RT-PCR) and cicular	USPAT;	2003/02/11 12:38
			US-PGPUB;	
			DERWENT	
4	405	(mRNA near5 RT-PCR) and circular	USPAT;	2003/02/11 12:38
			US-PGPUB;	
			DERWENT	/ /
5	332	((mRNA near5 RT-PCR) and circular) and	USPAT;	2003/02/11 12:39
		ligat\$	US-PGPUB;	
1 _			DERWENT	
6	62	(((mRNA near5 RT-PCR) and circular) and	USPAT;	2003/02/11 12:44
		ligat\$) and primer adj1 extension	US-PGPUB;	
1 _		(num	DERWENT	0000 (00 (11 10 50
7	205	(mRNA near5 RT-PCR) and RNase adj1 H	USPAT;	2003/02/11 12:50
			US-PGPUB;	
8	21	-i	DERWENT	2002/02/11 12:21
8	21	circular adj1 cDNA	USPAT;	2003/02/11 13:31
			US-PGPUB; DERWENT	
9	409	Reverse adj1 transcriptase near5 RNase	USPAT:	2003/02/11 13:32
9	409	adj1 H	US-PGPUB;	2003/02/11 13:32
		1	DERWENT	
10	5	(Reverse adj1 transcriptase near5 RNase	USPAT:	2003/02/11 13:32
		adj1 H) and circular adj1 cDNA	US-PGPUB;	2003/02/11 13.32
		daji ii, and circular aaji comi	DERWENT	
	L		DETAILE	

Page 1









Related Resources improve the effectiveness of the technique. Based on first-hand experience with RACE, a critical review of numerous published variations of the key steps in the RACE method is presented. Also included is a detailed, effective protocol based on RNA ligase-mediated RACE/reverse ligation-mediated PCR, as well as a demonstration of its utility.

Publication Types:

- Review
- Review, Tutorial

PMID: 7573945 [PubMed - indexed for MEDLINE]




Write to the Help Desk NCBI | NLM | NIH Department of Health & Human Services Freedom of Information Act | Disclaimer

i686-pc-lmux-gnu Feb 4 2003 11.11:49









Entrez PubMed ☐1: Curr Issues Mol Biol 1999;1(1-2):13-20

Related Articles, Links

Synthesis of infectious viroids and other circular RNAs.

Rezaian MA.

PubMed Services CSIRO Plant Industry Horticulture Unit, Adelaide Laboratory, PO Box 350, Glen Osmond, SA 5064, Australia.

Viroids are small autonomously replicating RNAs that share structural features with other subviral circular single-stranded RNAs of plants. Viroids and other circular single-stranded RNAs can be synthesised in vitro by a PCR-based procedure using a simple set of reactions. Two end-to-end primers are selected from a desired region of the viroid, one for the synthesis of the first strand cDNA and another for the production of the second strand DNA. The second primer contains an 18 nucleotide T7 promoter at its 5' end, and is selected such that the G nucleotide at the transcription start site represents a G in the viroid. Linked reverse transcription-PCR results in linear double-stranded DNA consisting of the viroid sequence and the T7 promoter. Run-off transcription of the PCR product allows the synthesis of exact-length linear viroid RNA which can be circularised by T4 RNA ligase following an enzymic modification of the 5' triphosphate to a monophosphate. This procedure results in authentic viroid molecules and obviates the need for construction and cloning of DNA in the form of tandem repeats for infectivity tests. It also allows PCR-based manipulation of circular RNAs, thus greatly simplifying structure-function analyses of viroid molecules.

Related Resources

PMID: 11475697 [PubMed - indexed for MEDLINE]



Write to the Help Desk
NCBI | NLM | NIH
Department of Health & Human Services
Freedom of Information Act | Disclaimer

i686-pe-linux-gnu Feb 4 2003 11:11:49

